

Discovery of the spectroscopic binary nature of three bright southern Cepheids

L. Szabados¹, R. I. Anderson², A. Derekas^{1,3}, L. L. Kiss^{1,3,4}, T. Szalai⁵,
P. Székely⁶, J. L. Christiansen⁷

¹*Konkoly Observatory, Research Centre for Astronomy and Earth Sciences, Hungarian Academy of Sciences, H-1121 Budapest, Konkoly Thege Miklós út 15-17, Hungary*

²*Observatoire de Genève, Université de Genève, 51 Ch. des Maillettes, CH-1290 Versoix, Switzerland*

³*Sydney Institute for Astronomy, School of Physics, University of Sydney, NSW 2006, Australia*

⁴*ELTE Gothard-Lendület Research Group, H-9700 Szombathely, Szent Imre herceg út 112, Hungary*

⁵*Department of Optics and Quantum Electronics, University of Szeged, Dóm tér 9, H-6720 Szeged, Hungary*

⁶*Department of Experimental Physics, University of Szeged, Szeged H-6720, Hungary*

⁷*SETI Institute/NASA Ames Research Center, M/S 244-30, Moffett Field, CA 94035, USA*

Accepted Received ; in original form

ABSTRACT

We present the analysis of spectroscopic radial velocity and photometric data of three bright Galactic Cepheids: LR Trianguli Australis, RZ Velorum, and BG Velorum. Based on new radial velocity data, these Cepheids have been found to be members in spectroscopic binary systems.

The ratio of amplitudes of peak-to-peak radial velocity amplitude and photometric amplitude indicates the presence of a companion for LR TrA and BG Vel. *IUE* spectra indicate that the companions of RZ Vel and BG Vel cannot be hot stars.

The analysis of all available photometric data revealed that the pulsation period of RZ Vel and BG Vel varies monotonically, due to stellar evolution. Moreover, the longest period Cepheid in this sample, RZ Vel, shows period fluctuations superimposed on the monotonic period increase. The light-time effect interpretation of the observed pattern needs long-term photometric monitoring of this Cepheid. The pulsation period of LR TrA has remained constant since the discovery of its brightness variation.

Using statistical data, it is also shown that a large number of spectroscopic binaries remains to be discovered among bright classical Cepheids.

Key words: stars: variables: Cepheids – binaries: spectroscopic

1 INTRODUCTION

Classical Cepheid variable stars are primary distance indicators and rank among standard candles for establishing the cosmic distance scale, owing to the famous period-luminosity (P - L) relationship. Companions to Cepheids, however, complicate the situation. The contribution of the secondary star to the observed brightness has to be taken into account when involving any particular Cepheid in the calibration of the P - L relationship.

Binaries among Cepheids are not rare at all: their frequency exceeds 50 per cent for the brightest Cepheids, while among the fainter Cepheids an observational selection effect encumbers revealing binarity (Szabados 2003a).

Owing to some observational projects aimed at obtaining new radial velocities (RVs) of numerous Cepheids carried out during the last decades, a part of the selection effect has been removed. This progress is visualized in Fig. 1 where the current situation is compared with that 20 years ago. The data have been taken

from the on-line data base on binaries among Galactic Cepheids (<http://www.konkoly.hu/CEP/orbit.html>). To get rid of the fluctuation at the left part of the diagram, brightest Cepheids ($\langle V \rangle < 5$ mag) were merged in a single bin because such stars are extremely rare among Cepheids – see the histogram in Fig. 2.

In the case of pulsating variables, like Cepheids, spectroscopic binarity manifests itself in a periodic variation of the γ -velocity (i.e., the RV of the mass centre of the Cepheid). In practice, the orbital RV variation of the Cepheid component is superimposed on the RV variations of pulsational origin. To separate orbital and pulsational effects, knowledge of the accurate pulsation period is essential, especially when comparing RV data obtained at widely differing epochs. Therefore, the pulsation period and its variations have been determined with the method of the $O - C$ diagram (Sterken 2005) for each target Cepheid. Use of the accurate pulsation period obtained from the photometric data is a guarantee for the correct phase matching of the (usually less precise) RV data.

In this paper we point out spectroscopic binarity of three




Figure 1. Percentage of known binaries among Galactic classical Cepheids as a function of the mean apparent visual brightness in 1993 and 2013. The decreasing influence of the observational selection effect is noticeable.




Figure 2. Histogram showing the number distribution of known Galactic classical Cepheids as a function of their mean apparent visual brightness.

bright Galactic Cepheids by analysing RV data. The structure of this paper is as follows. The new observations and the equipment utilized are described in Sect. 2. Section 3 is devoted to the results on the three new spectroscopic binary (SB) Cepheids: LR Trianguli Australis, RZ Velorum, and BG Velorum. Basic information on these Cepheids are found in Table 1. Finally, Sect. 4 contains our conclusions.

2 NEW OBSERVATIONS

2.1 Spectra from Siding Spring Observatory

We performed an RV survey of Cepheids with the 2.3 m ANU telescope located at Siding Spring Observatory, Australia. The main aim of the project was to detect Cepheids in binary systems by measuring changes in the mean values of their RV curve which can be

Table 1. Basic data of the programme stars and the number of spectra.

Cepheid	$\langle V \rangle$ mag	P (d)	Mode of pulsation	No. of spectra SSO	<i>CORALIE</i>
LR TrA	7.80	2.428289	first overtone	10	32
RZ Vel	7.13	20.398532	fundamental	30	67
BG Vel	7.69	6.923843	fundamental	27	33

interpreted as the orbital motion of the Cepheid around the centre of mass in a binary system (change of γ -velocity). The target list was compiled to include Cepheids with single-epoch RV phase curve or without any published RV data. Several Cepheids suspected members in SB systems were also put on the target list. On 64 nights between October 2004 and March 2006 we monitored 40 Cepheids with pulsation periods between 2 and 30 days. Additional spectra of some targets were obtained in August 2007.

Medium-resolution spectra were taken with the Double Beam Spectrograph using the 1200 mm^{-1} gratings in both arms of the spectrograph. The projected slit width was $2''$ on the sky, which was about the median seeing during our observations. The spectra covered the wavelength ranges $4200\text{--}5200 \text{ \AA}$ in the blue arm and $5700\text{--}6700 \text{ \AA}$ in the red arm. The dispersion was 0.55 \AA px^{-1} , leading to a nominal resolution of about 1 \AA .

All spectra were reduced with standard tasks in IRAF¹. Reduction consisted of bias and flat field corrections, aperture extraction, wavelength calibration, and continuum normalisation. We checked the consistency of wavelength calibrations via the constant positions of strong telluric features, which proved the stability of the system. Radial velocities were determined only for the red arm data with the task *fxcor*, applying the cross-correlation method using a well-matching theoretical template spectrum from the extensive spectral library of Munari et al. (2005). Then, we made barycentric corrections to every single RV value. This method resulted in a $1\text{--}2 \text{ km s}^{-1}$ uncertainty in the individual RVs, while further tests have shown that our absolute velocity frame was stable to within $\pm 2\text{--}3 \text{ km s}^{-1}$. This level of precision is sufficient to detect a number of Cepheid companions, as they can often cause γ -velocity changes well above 10 km s^{-1} .

Discovery of six SBs among the 40 target Cepheids was already reported by Szabados et al. (2013). Binarity of the three Cepheids announced here could be revealed by involving independently obtained additional data (see Sect. 2.2). The individual RV data of the rest of the Cepheid targets will be published together with the results of the analysis of the spectra.

2.2 CORALIE observations from La Silla

All three Cepheids were among the targets during multiple observing campaigns between April 2011 and May 2012 using the fiber-fed high-resolution ($R \sim 60000$) echelle spectrograph *CORALIE* mounted at the Swiss 1.2 m Euler telescope at ESO La Silla Observatory, Chile. The instrument's design is described in Queloz et al. (2001); recent instrumental updates are found in Ségransan et al. (2010).

When it turned out that these three Cepheids have variable γ -velocities, several new spectra were obtained in December 2012 - January 2013 and April 2013.

The spectra are reduced by the efficient online reduction pipeline that performs bias correction, cosmics removal, and flat-fielding using tungsten lamps. ThAr lamps are used for the wavelength calibration. The reduction pipeline directly determines the RV via cross-correlation (Baranne et al. 1996) using a mask that resembles a G2 spectral type. The RV stability of the instrument is excellent and for non-pulsating stars the RV precision is limited

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

by photon-noise, see e.g., Pepe et al. (2002). However, the precision achieved for Cepheids is lower due to line asymmetries. We estimate a typical precision of $\sim 0.1 \text{ km s}^{-1}$ (including systematics due to pulsation) per data point for our data.

3 RESULTS FOR INDIVIDUAL CEPHEIDS

3.1 LR Trianguli Australis

Accurate value of the pulsation period The brightness variability of LR TrA (HD 137626, $\langle V \rangle = 7.80$ mag) was revealed by Strohmeier et al. (1966) based on the Bamberg photographic patrol plates. The Cepheid nature of variability and the first values of the pulsation period was determined by Eggen (1983). This Cepheid pulsates in the first overtone mode, therefore it has a small pulsational amplitude and nearly sinusoidal light and velocity curves.

In the case of Cepheids pulsating with a low amplitude, the $O - C$ diagram constructed for the median brightness (the midpoint between the faintest and the brightest states) is more reliable than that based on the moments of photometric maxima (Derekas et al. 2012). Therefore we determined the accurate value of the pulsation period by constructing an $O - C$ diagram for the moments of median brightness on the ascending branch of light curve since this is the phase when the brightness variations are steepest during the whole pulsational cycle.

All published photometric observations of LR TrA covering three decades were re-analysed in a homogeneous manner to determine seasonal moments of the chosen light curve feature. The relevant data listed in Table 2 are as follows:

Column 1: Heliocentric moment of the selected light curve feature (median brightness on the ascending branch for LR TrA, maximum brightness for both RZ Vel and BG Vel, see Tables 6 and 10, respectively);

Col. 2: Epoch number, E , as calculated from Eq. 1:

$$C = 2453104.9265 + 2.428289 \times E \pm 0.0037 \pm 0.000003 \quad (1)$$

(this ephemeris has been obtained by the weighted least squares parabolic fit to the $O - C$ differences);

Col. 3: the corresponding $O - C$ value;

Col. 4: weight assigned to the $O - C$ value (1, 2, or 3 depending on the quality of the light curve leading to the given difference);

Col. 5: reference to the origin of data.

The $O - C$ diagram of LR TrA based on the $O - C$ values listed in Table 2 is plotted in Fig. 3. The plot can be approximated by a constant period by the ephemeris (1) for the moments of median brightness on the ascending branch. The scatter of the points in Fig. 3 reflects the observational error and uncertainties in the analysis of the data.

Binarity of LR TrA There are no earlier RV data on this bright Cepheid. Our new data listed in Tables 3 and 4 have been folded on the accurate pulsation period given in the ephemeris (see Eq. 1). The merged RV phase curve is plotted in Fig. 4. Both individual data series could be split into seasonal subsets.

Variability in the γ -velocity is obvious. The γ -velocities (together with their uncertainties) are listed in Table 5. The γ -velocity in 2007 is more uncertain than in other years because this value is based on a single spectrum. Systematic errors can be excluded. Dozens of Cepheids in our sample with non-varying γ -velocities indicate stability of the equipment and reliability of the data reduction. Figure 5 is a better visualisation of the temporal variation in

Table 2. $O - C$ values of LR Trianguli Australis (see the description in Sect. 3.1).

JD _⊙ 2400000 +	E	$O - C$	W	Data source
45018.7822	-3330	0.0581	3	Eggen (1983)
47633.9607	-2253	-0.0307	3	Antonello et al. (1990)
47939.9568	-2127	0.0010	2	<i>Hipparcos</i> (ESA 1997)
48139.0426	-2045	-0.0329	3	<i>Hipparcos</i> (ESA 1997)
48440.1554	-1921	-0.0279	3	<i>Hipparcos</i> (ESA 1997)
48750.9547	-1793	-0.0496	3	<i>Hipparcos</i> (ESA 1997)
49814.6064	-1355	0.0115	3	Berdnikov (2008)
50370.7115	-1126	0.0384	3	Berdnikov (2008)
50574.6393	-1042	-0.0101	3	Berdnikov (2008)
50909.7531	-904	-0.0001	3	Berdnikov (2008)
51264.2883	-758	0.0049	3	Berdnikov (2008)
51650.4058	-599	0.0244	3	Berdnikov (2008)
51958.8010	-472	0.0269	2	Berdnikov (2008)
52041.3435	-438	0.0076	2	ASAS (Pojmanski 2002)
52366.7222	-304	-0.0044	3	Berdnikov (2008)
52500.2709	-249	-0.0116	3	ASAS (Pojmanski 2002)
52769.8038	-138	-0.0188	3	ASAS (Pojmanski 2002)
53102.5159	-1	0.0177	3	Berdnikov (2008)
53104.9151	0	-0.0114	3	ASAS (Pojmanski 2002)
53520.1818	171	0.0179	3	ASAS (Pojmanski 2002)
53840.7137	303	0.0156	3	ASAS (Pojmanski 2002)
54251.0850	472	0.0061	3	ASAS (Pojmanski 2002)
54615.3163	622	-0.0060	3	ASAS (Pojmanski 2002)
54960.1214	764	-0.0179	3	ASAS (Pojmanski 2002)




Figure 3. $O - C$ diagram of LR TrA. The plot can be approximated by a constant period.

Table 3. RV values of LR Trianguli Australis from the SSO spectra. This is only a portion of the full version available online only.

JD _⊙ 2400000 +	v_{rad} (km s^{-1})
53599.9325	-21.2
53600.9086	-32.0
53603.9327	-27.6
53605.9290	-31.0
53805.1657	-29.3




Figure 4. Merged RV phase curve of LR TrA. Different symbols mean data from different years: 2005: filled triangles; 2006: empty triangles; 2007: triangular star; 2012: filled circles; 2013: empty circles. Zero phase was arbitrarily chosen at JD 2400 000.0 (in all merged RV phase curves in this paper).




Figure 5. Temporal variation in the γ -velocity of LR TrA. The symbols for the different data sets are the same as in Fig. 4.

Table 4. CORALIE velocities of LR Trianguli Australis. This is only a portion of the full version available online only.

JD _☉ 2400 000 +	v_{rad} (km s ⁻¹)	σ (km s ⁻¹)
55938.8701	-27.97	0.05
55938.8718	-28.10	0.05
55939.8651	-29.85	0.02
55940.8686	-22.40	0.03
55941.8579	-33.14	0.04

Table 5. γ -velocities of LR Trianguli Australis.

Mid-JD 2400 000+	v_{γ} (km s ⁻¹)	σ (km s ⁻¹)	Data source
53603	-25.5	0.5	present paper
53808	-24.8	0.5	present paper
54331	-29.0	1.0	present paper
55981	-27.5	0.1	present paper
56344	-26.4	0.1	present paper

the γ -velocity. The seasonal drift in the γ -velocity is compatible with both short and long orbital periods.

The photometric contribution of the companion star decreases the observable amplitude of the brightness variability as deduced from the enhanced value of the ratio of the RV and photometric amplitudes (Klagyivik & Szabados 2009). This is an additional (**although slight**) indication of binarity of LR TrA.

3.2 RZ Velorum

Accurate value of the pulsation period The brightness variability of RZ Vel (HD 73502, $\langle V \rangle = 7.13$ mag) was revealed by Cannon (Pickering 1909). The Cepheid nature of variability and the pulsation period was established by Hertzsprung (1936) based on the Harvard and Johannesburg photographic plate collection which was further investigated by Oosterhoff (1936).

This is the longest period Cepheid in this paper and it has been frequently observed from the 1950s, first photoelectrically, then in the last decades by CCD photometry. The photometric coverage of RZ Vel was almost continuous in the last twenty years thanks to observational campaigns by Berdnikov (2008) and his co-workers, as well as the ASAS photometry (Pojmanski 2002).

Long-period Cepheids are usually fundamental pulsators and they oscillate with a large amplitude resulting in a light curve with sharp maximum.

The $O - C$ diagram of RZ Vel was constructed for the moments of maximum brightness based on the photoelectric and CCD photometric data (see Table 6). The weighted least squares parabolic fit to the $O - C$ values resulted in the ephemeris:

$$C = 2442453.6630 + 20.398532 \times E + 1.397 \times 10^{-6} E^2 \quad (2)$$

$$\pm 0.0263 \quad \pm 0.000080 \quad \pm 0.191 \times 10^{-6}$$

The $O - C$ diagram of RZ Vel plotted in Fig. 6 indicates a continuously increasing pulsation period with a period jitter superimposed. This secular period increase has been caused by stellar evolution: while the Cepheid crosses the instability region towards lower temperatures in the Hertzsprung-Russell diagram, its pulsation period is increasing. Continuous period variations (of either sign) often occur in the pulsation of long-period Cepheids (Szabados 1983).

Figure 7 shows the $O - C$ residuals after subtracting the parabolic fit defined by Eq. 2. If the wave-like fluctuation seen in this $\Delta(O - C)$ diagram turns out to be periodic, it would correspond to a light-time effect in a binary system. In line with the recent shortening in the pulsation period, the current value of the pulsation period is 20.396671 ± 0.000200 days (after JD 2452 300).

Binarity of RZ Vel There are several data sets of RV observations available in the literature for RZ Vel: those published by Stibbs (1955), Lloyd Evans (1968, 1980), Coulson & Caldwell (1985), Bersier (2002), and Nardetto et al. (2006). Our individual RV data are listed in Tables 7 and 8.

Based on these data, the RV phase curve has been constructed using the 20.398532 d pulsation period appearing in Eq. 2. In view of the complicated pattern of the $O - C$ diagram the RV data have been folded on by taking into account the proper phase correction for different data series. The merged RV phase curve is plotted in Fig. 8. For the sake of clarity, RV data obtained before JD 2450 000 have not been plotted here because of the wider scatter of these early RV data but the γ -velocities were determined for each data set. The individual data series are denoted with different symbols: filled squares mean data by Bersier (2002), empty squares those by Nardetto et al. (2006), our 2005, 2006, 2012, 2013 data are denoted

Table 6. $O - C$ values of RZ Velorum (description of the columns is given in Sect. 3.1).

JD_{\odot} 2 400 000 +	E	$O - C$	W	Data source
33784.5646	-425	0.2777	1	Eggen et al. (1957)
34804.5174	-375	0.3039	1	Walraven et al. (1958)
34845.2119	-373	0.2013	3	Eggen et al. (1957)
35192.0024	-356	0.2168	1	Irwin (1961)
40760.8647	-83	0.2799	3	Pel (1976)
41719.0924	-36	-0.2234	3	Madore (1975)
41862.1249	-29	0.0193	3	Dean et al. (1977)
42453.6330	0	-0.0030	3	Dean et al. (1977)
44371.0472	94	-0.0778	3	Coulson & Caldwell (1985)
44391.3842	95	-0.1393	2	Eggen (1982)
45003.2906	125	-0.1889	3	Coulson & Caldwell (1985)
48226.4369	283	-0.0107	3	<i>Hipparcos</i> (ESA 1997)
48797.5877	311	-0.0188	3	<i>Hipparcos</i> (ESA 1997)
49185.1653	330	-0.0133	1	Walker & Williams (unpublished)
49817.8011	361	0.2680	3	Berdnikov (2008)
50144.1979	377	0.2883	2	Bersier (2002)
50389.0443	389	0.3524	3	Berdnikov (2008)
50511.3662	395	0.2831	3	Bersier (2002)
50572.4468	398	0.1681	3	Berdnikov (2008)
50899.0581	414	0.4029	3	Berdnikov (2008)
51266.1488	432	0.3200	3	Berdnikov (2008)
51653.7650	451	0.3641	3	Berdnikov (2008)
51939.2846	465	0.3042	2	ASAS (Pojmanski 2002)
51959.7692	466	0.3903	3	Berdnikov (2008)
52347.4262	485	0.4752	3	Berdnikov (2008)
52653.3896	500	0.4606	3	ASAS (Pojmanski 2002)
52653.4100	500	0.4810	3	Berdnikov (2008)
53000.1794	517	0.4754	3	ASAS (Pojmanski 2002)
53000.2610	517	0.5570	3	Berdnikov (2008)
53428.4384	538	0.3652	3	ASAS (Pojmanski 2002)
53754.8864	554	0.4367	3	ASAS (Pojmanski 2002)
54183.1657	575	0.3468	3	ASAS (Pojmanski 2002)
54509.5729	591	0.3775	3	ASAS (Pojmanski 2002)
54815.4343	606	0.2609	3	ASAS (Pojmanski 2002)
55121.3569	621	0.2055	2	ASAS (Pojmanski 2002)






Figure 6. $O - C$ diagram of RZ Vel. The plot can be approximated by a parabola indicating a continuously increasing period.

Figure 7. $\Delta(O - C)$ diagram of RZ Vel.

Figure 8. Radial velocity phase curve of RZ Vel. Data obtained between 1996 and 2013 are included in this plot. The meaning of various symbols is explained in the text.

Figure 9. γ -velocities of RZ Velorum. The symbols for the different data sets are the same as in Fig. 8.

Table 7. RV values of RZ Velorum from the SSO spectra. This is only a portion of the full version available online only.

JD_{\odot} 2 400 000 +	v_{rad} (km s^{-1})
53307.2698	4.2
53310.2504	1.4
53312.2073	9.0
53364.2062	49.6
53367.1823	27.5

Table 8. *CORALIE* velocities of RZ Velorum. This is only a portion of the full version available online only.

JD _⊙ 2 400 000 +	v_{rad} (km s ⁻¹)	σ (km s ⁻¹)
55654.5528	-3.08	0.02
55656.6626	5.23	0.01
55657.6721	9.86	0.02
55659.6585	18.85	0.03
55662.5137	31.50	0.01

Table 9. γ -velocities of RZ Velorum.

Mid-JD 2 400 000+	v_{γ} (km s ⁻¹)	σ (km s ⁻¹)	Data source
34009	25.5	1.5	Stibbs (1955)
40328	22.1	1.5	Lloyd Evans (1968, 1980)
42186	29.2	1.0	Coulson & Caldwell (1985)
44186	22.6	1.0	Coulson & Caldwell (1985)
44736	24.4	1.0	Coulson & Caldwell (1985)
50317	25.1	0.2	Bersier (2002)
53184	24.0	0.5	Nardetto et al. (2006)
53444	26.9	0.6	present paper
53783	28.8	1.0	present paper
55709	25.6	0.1	present paper
56038	25.3	0.1	present paper

by filled and empty triangles, filled and empty circles, respectively. The wide scatter in this merged RV phase curve plotted in Fig. 8 is due to a variable γ -velocity.

The γ -velocities determined from each data set (including the earlier ones) are listed in Table 9 and are plotted in Fig. 9. This latter plot implies that RZ Vel is really a spectroscopic binary as suspected by Bersier (2002) based on a much poorer observational material (before JD 2 450 500). An orbital period of about 5600-5700 days is compatible with the data pattern in both Fig. 7 and Fig. 9 but the phase relation between the light-time effect fit to the $\Delta(O - C)$ curve and the orbital radial velocity variation phase curve obtained with this formal period is not satisfactory.

3.3 BG Velorum

Accurate value of the pulsation period The brightness variability of BG Vel (HD 78801, $\langle V \rangle = 7.69$ mag) was revealed by Cannon (Pickering 1909). Much later O’Leary (1937) independently discovered its light variations but he also revealed the Cepheid nature and determined the pulsation period based on photographic plates obtained at the Riverview College Observatory. Van Houten (1950) also observed this Cepheid photographically in Johannesburg but these early data are unavailable, therefore we only mention their studies for historical reasons.

This Cepheid is a fundamental mode pulsator. The $O - C$ differences of BG Velorum calculated for brightness maxima are listed in Table 10. These values have been obtained by taking into account the constant and linear terms of the following weighted parabolic fit:


$$C = 2\,453\,031.4706 + 6.923\,843 \times E + 2.58 \times 10^{-8} E^2 \quad (3)$$

$$\pm 0.0020 \pm 0.000\,007 \quad \pm 0.27 \times 10^{-8}$$

The parabolic nature of the $O - C$ diagram, i.e., the continuous increase in the pulsation period is clearly seen in Fig. 10. This

Table 10. $O - C$ values of BG Velorum (description of the columns is given in Sect. 3.1).

JD _⊙ 2 400 000 +	E	$O - C$	W	Data source
34856.5526	-2625	0.1699	3	Walraven et al. (1958)
35237.3813	-2570	0.1872	3	Irwin (1961)
40748.6592	-1774	0.0861	3	Pel (1976)
42853.4433	-1470	0.0219	3	Dean (1977)
44300.5426	-1261	0.0380	3	Berdnikov (2008)
48136.3167	-707	0.0031	3	<i>Hipparcos</i> (ESA 1997)
48627.9239	-636	0.0174	3	<i>Hipparcos</i> (ESA 1997)
50379.6329	-383	-0.0058	3	Berdnikov (2008)
50573.4987	-355	-0.0076	3	Berdnikov (2008)
50905.8549	-307	0.0041	3	Berdnikov (2008)
51265.9127	-255	0.0221	3	Berdnikov (2008)
51646.7345	-200	0.0325	3	Berdnikov (2008)
51937.5210	-158	0.0176	3	ASAS (Pojmanski 2002)
51958.2712	-155	-0.0038	3	Berdnikov (2008)
52359.8640	-97	0.0062	3	ASAS (Pojmanski 2002)
52359.8778	-97	0.0200	3	Berdnikov (2008)
52650.6575	-55	-0.0017	3	Berdnikov (2008)
52726.8212	-44	-0.0003	3	ASAS (Pojmanski 2002)
53003.7916	-4	0.0164	3	Berdnikov (2008)
53031.4758	0	0.0052	3	ASAS (Pojmanski 2002)
53336.1201	44	0.0004	1	<i>INTEGRAL</i> OMC
53460.7390	62	-0.0099	3	ASAS (Pojmanski 2002)
53779.2202	108	-0.0254	3	ASAS (Pojmanski 2002)
54180.8337	166	0.0052	3	ASAS (Pojmanski 2002)
54540.8499	218	-0.0185	3	ASAS (Pojmanski 2002)
54838.5810	261	-0.0126	3	ASAS (Pojmanski 2002)
55143.2425	305	-0.0002	2	ASAS (Pojmanski 2002)

**Figure 10.** $O - C$ diagram of BG Vel. The plot can be approximated by a parabola indicating a continuously increasing pulsation period.

parabolic trend corresponds to a continuous period increase of $(5.16 \pm 0.54) \times 10^{-8}$ d/cycle, i.e., $\Delta P = 0.000272$ d/century. This tiny period increase has been also caused by stellar evolution as in the case of RZ Vel.

The fluctuations around the fitted parabola in Fig. 10 do not show any definite pattern: see the $\Delta(O - C)$ diagram in Fig. 11.

Binarity of BG Vel There are earlier RV data of this Cepheid obtained by Stibbs (1955) and Lloyd Evans (1980). Variability in the γ -velocity is seen in the merged phase diagram of all RV data of BG Velorum plotted in Fig. 12. In this diagram, our 2005-2006 data (listed in Table 11) are represented with empty circles, while 2012-2013 data (listed in Table 12) are denoted with filled circles, triangles represent Stibbs’ data, and \times symbols refer to Lloyd




Figure 11. $\Delta(O - C)$ diagram of BG Vel.




Figure 12. Merged RV phase curve of BG Vel. There is an obvious shift between the γ -velocities valid for the epoch of our data obtained in 2005-2006 and 2012-2013 (empty and filled circles, respectively). The other symbols are explained in the text.




Figure 13. γ -velocities of BG Velorum. The symbols for the different data sets are the same as in Fig. 12.

Table 11. RV values of BG Velorum from the SSO spectra. This is only a portion of the full version available online only.

JD _⊙ 2 400 000 +	v_{rad} (km s ⁻¹)
53312.2372	17.3
53364.2219	-0.2
53367.1992	20.5
53451.0000	20.0
53452.0021	23.8

Table 12. CORALIE velocities of BG Velorum. This is only a portion of the full version available online only.

JD _⊙ 2 400 000 +	v_{rad} (km s ⁻¹)	σ (km s ⁻¹)
55937.7555	24.13	0.02
55938.6241	7.77	0.02
55939.6522	-1.25	0.01
55941.6474	7.99	0.10
55942.6917	11.78	0.03

Table 13. γ -velocities of BG Velorum.

Mid-JD 2 400 000 +	v_{γ} (km s ⁻¹)	σ (km s ⁻¹)	Data source
34096	11.4	1.5	Stibbs (1955)
40545	8.4	1.5	Lloyd Evans (1980)
53572	12.6	0.6	present paper
56043	10.3	0.1	present paper

Evans' data. Our RV data have been folded with the period given in the ephemeris Eq. 3 omitting the quadratic term. Data obtained by Stibbs and Lloyd Evans have been phased with the same period but a proper correction has been applied to allow for the phase shift due to the parabolic $O - C$ graph.

The γ -velocities determined from the individual data sets are listed in Table 13 and plotted in Fig. 13. **Because** no annual shift is seen in the γ -velocities between two consecutive years (2005-2006 and 2012-2013), the orbital period cannot be short, probably it exceeds a thousand days.

Similarly to the case of LR TrA, BG Vel is also characterised by an excessive value for the ratio of RV and photometric amplitudes indicating the **possible** presence of a companion (see Fig. 14).

4 CONCLUSIONS

We pointed out that three bright southern Galactic Cepheids, LR Trianguli Australis, RZ Velorum, and BG Velorum have a variable γ -velocity implying their membership in SB systems. RV values of other target Cepheids observed with the same equipment in 2005-2006 and 2012 testify that this variability in the γ -velocity is not of instrumental origin, nor an artefact caused by the analysis.

The available RV data are insufficient to determine the orbital period and other elements of the orbits. However, some inferences can be made from the temporal variations of the γ -velocity. An orbital period of 5600-5700 days of the RZ Vel system is compatible with the data pattern. In the case of BG Vel, short orbital periodicity can be ruled out. For LR TrA, even the range of the possible orbital periods remains uncertain.

The value of the orbital period for spectroscopic binary systems involving a Cepheid component is often unknown: according to the on-line data base (Szabados 2003a) the orbital period has been determined for about 20% of the known SB Cepheids. The majority of known orbital periods exceeds a thousand days.

A companion star may have various effects on the observable photometric properties of the Cepheid component. Various pieces of evidence of duplicity based on the photometric criteria are discussed by Szabados (2003b) and Klagyivik & Szabados (2009). As




Figure 14. The slightly excessive value of the A_{VRAD}/A_B amplitude ratio of LR TrA and BG Vel (large circles) with respect to the average value characteristic at the given pulsation period is an independent indication of the presence of a companion star. This is a modified version of Fig. 4f of Klagyivik & Szabados (2009). For the meaning of various symbols, see the original figure.

to our targets, there is no obvious sign of a companion from optical multicolour photometry. This indicates that the companion star cannot be much hotter than any of the Cepheids discussed here. There is, however, a phenomenological parameter, viz. the ratio of RV and photometric amplitudes (Klagyivik & Szabados 2009) whose excessive value is a further hint at the probable existence of a companion for both LR TrA and BG Vel (see Fig. 14). Moreover, the *IUE* spectra of bright Cepheids analysed by Evans (1992) gave a constraint on the temperature of a companion to remain undetected in the ultraviolet spectra: in the case of RZ Vel, the spectral type of the companion cannot be earlier than A7, while for BG Vel this limiting spectral type is A0. Further spectroscopic observations are necessary to characterise these newly detected SB systems.

Our findings confirm the previous statement by Szabados (2003a) about the high percentage of binaries among classical Cepheids and the observational selection effect hindering the discovery of new cases (see also Fig. 1).

Regular monitoring of the radial velocities of a large number of Cepheids will be instrumental in finding more SBs among Cepheids. RV data to be obtained with the *Gaia* astrometric space probe (expected launch: September 2013) will certainly result in revealing new SBs among Cepheids brighter than 13-14th magnitude (Eyer et al. 2012). In this manner, the ‘missing’ SBs among Cepheids inferred from Fig. 1 can be successfully revealed within several years.

ACKNOWLEDGMENTS

This project has been supported by the ESTEC Contract No. 4000106398/12/NL/KML, the Hungarian OTKA Grants K76816, K83790, K104607, and MB08C 81013, as well as the European Community’s Seventh Framework Program (FP7/2007-2013) under grant agreement no. 269194, and the “Lendület-2009” Young Researchers Program of the Hungarian Academy of Sciences. AD was supported by the Hungarian Eötvös fellowship. AD has been supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. AD is very thankful to the staff at The Lodge in Siding Spring Observatory for their hospitality and the very nice food, making the time spent there lovely and special.

Part of the research leading to these results has received funding from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007–2013)/ERC grant agreement n°227224 (PROSPERITY). The *INTEGRAL* photometric data, pre-processed by ISDC, have been retrieved from the OMC Archive at CAB (INTA-CSIC). We are indebted to Stanley Walker for sending us some unpublished photoelectric observational data. **Our thanks are also due to the referee and Dr. Mária Kun for their critical remarks leading to a considerable improvement in the presentation of the results.**

REFERENCES

- Antonello E., Poretti E., Reduzzi L. 1990, *A&A*, 236, 138
 Baranne A. et al. 1996, *A&AS*, 119, 373
 Berdnikov L. N. 2008, *VizieR On-line Data Catalog: II/285*
 Bersier D. 2002, *ApJS*, 140, 465
 Coulson I. M., Caldwell J. A. R. 1985, *SAAO Circ.*, No. 9, 5
 Dean J. F. 1977, *MNASSA*, 36, 3
 Dean J. F., Cousins A. W. J., Bywater R. A., Warren P. R. 1977, *Mem RAS*, 83, 69
 Derekas A. et al. 2012, *MNRAS*, 425, 1312
 Eggen O. J. 1982, *ApJS*, 50, 199
 Eggen O. J. 1983, *AJ*, 88, 361
 Eggen O. J. 1985, *AJ*, 90, 1297
 Eggen O. J., Gascoigne S. C. B., Burr E. J. 1957, *MNRAS*, 117, 406
 ESA 1997, *The Hipparcos and Tycho Catalogues*, ESA SP-1200
 Evans N. R. 1992, *ApJ*, 384, 220
 Eyer L. et al. 2012, *Ap&SS*, 341, 207
 Hertzsprung E. 1936, *Bull. Astr. Inst. Neth.*, 8, 25
 Irwin J. B. 1961, *ApJS*, 6, 253
 Klagyivik P., Szabados L. 2009, *A&A*, 504, 959
 Lloyd Evans T. 1968, *MNRAS*, 141, 109
 Lloyd Evans T. 1980, *SAAO Circ.*, 1, No. 5, 257
 Madore B. F. 1975, *ApJS*, 29, 219
 Munari U., Sordo R., Castelli F., Zwitter T. 2005, *A&A*, 442, 1127
 Nardetto N., Mourard D., Kervella P., Mathias Ph., Mérand A., Bersier D. 2006, *A&A*, 453, 309
 O’Leary W. 1937, *Riverview Coll. Obs. Publications*, No. 4, 49
 Oosterhoff P. Th. 1936, *Bull. Astr. Inst. Neth.*, 8, 29
 Pel J. W. 1976, *A&AS*, 24, 413
 Pepe F., Mayor M., Galland F., Naef D., Queloz D., Santos N. C., Udry S., Burnet M. 2002, *A&A*, 388, 632
 Pickering E. C. 1909, *Harvard Coll. Obs. Circ.*, No. 151, 1
 Pojmanski G. 2002, *AcA*, 52, 397
 Queloz D. et al. 2001, *Messenger*, No. 105, 1
 Ségransan D. et al. 2010, *A&A*, 511, 45
 Sterken C. 2005, in Sterken C. ed., *The Light-Time Effect in Astrophysics*, ASP Conf. Ser. 335, Astron. Soc. Pac., San Francisco, p. 3
 Stibbs D. W. N. 1955, *MNRAS*, 115, 363
 Strohmeier W., Fischer H., Ott H. 1966, *Inf. Bull. Var. Stars*, No. 120, 1
 Szabados L. 1983, *Ap&SS*, 96, 185
 Szabados L. 2003a, *Inf. Bull. Var. Stars*, 5394
 Szabados L. 2003b, in *Recent Res. Devel. Astron. & Astrophys.*, 1, (Trivandrum: Research Signpost), p. 787
Szabados L. et al. 2013, *MNRAS*, 430, 2018
 van Houten C. J. 1977, *Leiden Ann.*, 20, 223

Walraven Th., Muller A. B., Oosterhoff P. T. 1958, Bull. Astr. Inst. Neth., 14, 81

This paper has been typeset from a \TeX / \LaTeX file prepared by the author.